


In re: Nikolai Vladimirovich Bovin et al.  
Serial No.: To Be Assigned  
Filed: Concurrently Herewith  
Page 16 of 30

### REMARKS

The claims have been amended to better conform with U.S. practice, such as removing multiple dependencies. Applicants respectfully request substantive examination on the merits.

Respectfully submitted,



Jarett K. Abramson  
Registration No. 47,376  
Attorney for Applicants



20792

PATENT TRADEMARK OFFICE

### CERTIFICATE OF EXPRESS MAILING

Express Mail Label No. EL 920742783 US

Date of Deposit: December 21, 2001

I hereby certify that this correspondence is being deposited with the United States Postal Service "Express Mail Post Office to Addressee" service under 37 CFR 1.10 on the date indicated above and is addressed to: BOX PCT, Attn: DO/EQ/US, Commissioner for Patents, Washington, DC 20231.



Susan E. Freedman

Date of Signature: December 21, 2001

**VERSION WITH MARKINGS TO SHOW CHANGES MADE**

The following is an addendum to the concurrently filed Preliminary Amendment in the above-referenced application. This addendum includes a marked-up version of the changes made to the claims by the present Preliminary Amendment.

**In the Claims:**

1. (Amended) [Compound] A compound of the general formula (I)



wherein

X is an m-valent unit and

B are identical or different and denote K-R,

wherein

K is a bond or is  $A^1-(A^2-A^3)_k$ -sp, wherein

$A^1$  is  $(CH_2)_tY(CH_2)_u$ , wherein

Y is  $>C=O$ ,  $>NH$ ,  $-O-$ ,  $-S-$  or a bond,

t is an integer from 0 to 6 and

u is an integer from 0 to 6,

$A^2$  is  $-NHCO-$ ,  $-CONH-$ ,  $-OCONH-$  or  $SCONH-$ , or is  $-CO-$ ,

$A^3$  is  $(CH_2)_r$ ,  $O(CH_2)_r$ ,  $NH(CH_2)_r$ ,  $S(CH_2)_r$  or  $-(CHQ)-$ , wherein

r is an integer from 1 to 6 and

Q is a substituted or unsubstituted alkyl or aryl group,

sp is a divalent spacer or a bond, and

k is an integer from 5 to 100, and

R is hydrogen; a ligand suitable for specific bonding to a receptor;  
a marker molecule; or a catalytically active group; and

m is at least 2,

with the proviso that

- (1) in the compound at least one R is not hydrogen,
- (2) there are at least two K that are not a bond, and

In re: Nikolai Vladimirovich Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 18 of 30

- (3) X, B and m are so selected that an intermolecular association of the K in liquid phase by the formation of hydrogen bonds is possible, with formation of aggregates that present on the surface a plurality of R that are not hydrogen, and
- (4) the molar mass of the fragment  $X(K)_m$  is less than 20,000.

2. (Amended) [Compound] A compound according to claim 1, wherein the molar mass of the fragment  $X(K)_m$  is less than 4,000.

3. (Amended) [Compound] A compound according to [either] claim 1 [or claim 2], wherein

m is an integer from 2 to 4, and

X is  $CH_{4-m}$ ,  $NH_{3-m}$ ,  $N^+H_{4-m}$ ,  $>P^-$  (when  $m = 3$ ),  $>P^+<$  (when  $m = 4$ ),  $>B^-$  (when  $m = 3$ ), a linear atom group  $C_2H_{6-m}$ ,  $>CH(CH_2)_zCH<$ ,  $>C=C<$ ,  $>N-N<$ ,  $>N(CH_2)_zN<$  wherein  $z = 2 - 6$ , when  $m = 4$ , a carbocyclic atom group  $C_6H_6-m$ ,  $C_6H_{12-m}$ , or a heterocyclic atom group  $C_3N_3$  (when  $m = 3$ ),  $C_4N_2$  (when  $m = 4$ ).

4. (Amended) [Compound] A compound according to [any one of claims 1 to 3] claim 1, wherein there are at least 3 K.

5. (Amended) [Compound] A compound according to [any one of claims 1 to 4] claim 1, wherein at least two R are not hydrogen.

6. (Amended) [Compound] A compound according to [any one of claims 1 to 4] claim 1, wherein at least three R are not hydrogen.

7. (Amended) [Compound] A compound according to [any one of claims 1 to 6] claim 1, wherein the ligand R is a mono- or oligo-saccharide, a peptide, a mono- or oligo-nucleotide or a nucleic base and their derivatives and mimetics.

8. (Amended) [Compound] A compound according to claim 7, wherein the saccharide R is sialic acid, sialyl lactose, sialyl lactosamine, lactose, mannose,  $Gal\alpha 1-3Gal$ ,

In re: Nikolai Vladimi h Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 19 of 30

Gal $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal, GalNAc $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal, Neu5Ac $\alpha$ 2-6GalNAc, SiaLe<sup>A</sup>, SiaLe<sup>X</sup>, HSO<sub>3</sub>Le<sup>A</sup>, HSO<sub>3</sub>Le<sup>X</sup>, Gal $\alpha$ 1-3Gal $\beta$ 1-4GlcNAc, Gal $\alpha$ 1-3Gal $\beta$ 1-4Glc, HSO<sub>3</sub>GlcA $\beta$ 1-3Gal $\beta$ 1-4GlcNAc, N-acetyl-lactosamine or polylactosamine, or wherein the saccharide is sialic acid benzyl glycoside, HSO<sub>3</sub>GlcA $\beta$ 1-3Gal, HSO<sub>3</sub>GlcA $\beta$ 1-3Gal $\beta$ 1-4GlcNAc $\beta$ 1-3Gal $\beta$ 1-4Glc, GalNAc $\alpha$ , GalNAc $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal $\beta$ 1-4GlcNAc, Gal $\alpha$ 1-3(Fuc $\alpha$ 1-2)Gal $\beta$ 1-4GlcNAc, HSO<sub>3</sub>(Sia)Le<sup>X</sup>, HSO<sub>3</sub>(Sia)Le<sup>A</sup>, Le<sup>Y</sup>, GlcNAc $\beta$ 1-6(GlcNAc $\beta$ 1-3)Gal $\beta$ 1-4Glc, GalNAc $\beta$ 1-4(Neu5Ac $\alpha$ 2-3)Gal $\beta$ 1-4Glc, mannose-6-phosphate, GalNAc $\beta$ 1-4GlcNAc, oligo-sialic acid, N-glycolylneuraminic acid, Gal $\alpha$ 1-4Gal $\beta$ 1-4Glc, Gal $\alpha$ 1-4Gal $\beta$ 1-4GlcNAc.

9. (Amended) [Compound] A compound according to [any one of claims 1 to 8] claim 1, wherein

m is an integer from 2 to 4,  
 X is CH<sub>4-m</sub>,  
 A<sup>1</sup> is CH<sub>2</sub>,  
 A<sup>2</sup> is NHCO,  
 A<sup>3</sup> is CH<sub>2</sub>,  
 k is 8,  
 sp is (CH<sub>2</sub>)<sub>3</sub>CONHCH<sub>2</sub>CONHC<sub>6</sub>H<sub>4</sub>-4-CH<sub>2</sub>O- and  
 R is Neu5Ac $\alpha$ 2-6Gal $\beta$ 1-4GlcNAc.

10. (Amended) [Aggregate] An aggregate of the general formula (II):



wherein

X(B)<sub>m</sub> may be identical or different and denote a compound of the general formula (I), [as defined in any one of claims 1 to 11]



wherein

X is an m-valent unit and

B are identical or different and denote K-R,

wherein

K is a bond or is A<sup>1</sup>-(A<sup>2</sup>-A<sup>3</sup>)<sub>k</sub>-sp, wherein

In re: Nikolai Vladimirovich Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 20 of 30

$A^1$  is  $(CH_2)_t Y (CH_2)_u$ , wherein  
 $Y$  is  $>C=O$ ,  $>NH$ ,  $-O-$ ,  $-S-$  or a bond,  
 $t$  is an integer from 0 to 6 and  
 $u$  is an integer from 0 to 6,  
 $A^2$  is  $-NHCO-$ ,  $-CONH-$ ,  $-OCONH-$  or  $SCONH-$ , or is  $-CO-$ ,  
 $A^3$  is  $(CH_2)_r$ ,  $O(CH_2)_r$ ,  $NH(CH_2)_r$ ,  $S(CH_2)_r$  or  $-(CHQ)-$ , wherein  
 $r$  is an integer from 1 to 6 and  
 $Q$  is a substituted or unsubstituted alkyl or aryl group,  
 $sp$  is a divalent spacer or a bond, and  
 $k$  is an integer from 5 to 100, and  
 $R$  is hydrogen; a ligand suitable for specific bonding to a receptor;  
 a marker molecule; or a catalytically active group; and  
 $m$  is at least 2,  
 with the proviso that  
 (1) in the compound at least one  $R$  is not hydrogen,  
 (2) there are at least two  $K$  that are not a bond, and  
 (3)  $X$ ,  $B$  and  $m$  are so selected that an intermolecular association of the  $K$  in liquid phase  
 by the formation of hydrogen bonds is possible, with formation of aggregates that  
 present on the surface a plurality of  $R$  that are not hydrogen, and  
 (4) the molar mass of the fragment  $X(K)_m$  is less than 20,000, and  
 $n$  is from 2 to 100,000,  
 and wherein  $X(B)_m$  are non-covalently bonded.

11. (Amended) [Aggregate] An aggregate according to claim 10 having a leaf-like, linear, cyclic, polycyclic, polyhedral, spherical or dendritic structure.

12. (Amended) [Aggregate] An aggregate according to claim 10 [or 11] of two or more different compounds [according to any one of claims 1 to 9] comprising a compound of the general formula (I)



wherein

$X$  is an m-valent unit and

In re: Nikolai Vladimirovich Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 21 of 30

B are identical or different and denote K-R,

wherein

K is a bond or is  $A^1-(A^2-A^3)_k\text{-sp}$ , wherein

$A^1$  is  $(\text{CH}_2)_t\text{Y}(\text{CH}_2)_u$ , wherein

Y is  $>\text{C}=\text{O}$ ,  $>\text{NH}$ ,  $-\text{O}-$ ,  $-\text{S}-$  or a bond,

t is an integer from 0 to 6 and

u is an integer from 0 to 6,

$A^2$  is  $-\text{NHCO}-$ ,  $-\text{CONH}-$ ,  $-\text{OCONH}-$  or  $\text{SCONH}-$ , or is  $-\text{CO}-$ ,

$A^3$  is  $(\text{CH}_2)_r$ ,  $\text{O}(\text{CH}_2)_r$ ,  $\text{NH}(\text{CH}_2)_r$ ,  $\text{S}(\text{CH}_2)_r$  or  $-(\text{CHQ})-$ , wherein

r is an integer from 1 to 6 and

Q is a substituted or unsubstituted alkyl or aryl group,

sp is a divalent spacer or a bond, and

k is an integer from 5 to 100, and

R is hydrogen; a ligand suitable for specific bonding to a receptor;

a marker molecule; or a catalytically active group; and

m is at least 2,

with the proviso that

(1) in the compound at least one R is not hydrogen,

(2) there are at least two K that are not a bond, and

(3) X, B and m are so selected that an intermolecular association of the K in liquid phase by the formation of hydrogen bonds is possible, with formation of aggregates that present on the surface a plurality of R that are not hydrogen, and

(4) the molar mass of the fragment  $\text{X}(\text{K})_m$  is less than 20,000.

Claim 13 has been canceled.

14. (Amended) [Process] A method according to claim [13] 27, further comprising [in the case of a solution of the compound addition of] adding a concentrated salt solution, [a change in] changing the pH or the temperature, or [addition of] adding organic solvents.

In re: Nikolai Vladimirovich Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 22 of 30

15. (Amended) [Process] A method for changing the structure of [the] an aggregate [as defined in any one of claims 10 to 12, which comprises addition of] of the general formula (II)



wherein

X(B)<sub>m</sub> may be identical or different and denote a compound of the general formula (I),



wherein

X is an m-valent unit and

B are identical or different and denote K-R,

wherein

K is a bond or is  $A^1-(A^2-A^3)_k\text{-sp}$ , wherein

A<sup>1</sup> is  $(CH_2)_tY(CH_2)_u$ , wherein

Y is  $>C=O$ ,  $>NH$ ,  $-O-$ ,  $-S-$  or a bond,

t is an integer from 0 to 6 and

u is an integer from 0 to 6,

A<sup>2</sup> is  $-NHCO-$ ,  $-CONH-$ ,  $-OCONH-$  or  $SCONH-$ , or is  $-CO-$ ,

A<sup>3</sup> is  $(CH_2)_r$ ,  $O(CH_2)_r$ ,  $NH(CH_2)_r$ ,  $S(CH_2)_r$  or  $-(CHQ)-$ , wherein

r is an integer from 1 to 6 and

Q is a substituted or unsubstituted alkyl or aryl group,

sp is a divalent spacer or a bond, and

k is an integer from 5 to 100, and

R is hydrogen; a ligand suitable for specific bonding to a receptor;

a marker molecule; or a catalytically active group; and

m is at least 2,

with the proviso that

(1) in the compound at least one R is not hydrogen,

(2) there are at least two K that are not a bond, and

(3) X, B and m are so selected that an intermolecular association of the K in liquid phase by the formation of hydrogen bonds is possible, with formation of aggregates that present on the surface a plurality of R that are not hydrogen, and

(4) the molar mass of the fragment X(K)<sub>m</sub> is less than 20,000, and

In re: Nikolai Vladimirovich Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 23 of 30

n is from 2 to 100,000,  
and wherein  $X(B)_m$  are non-covalently bonded,  
further comprising adding a concentrated salt solution, [a change in] changing the  
temperature or the pH [or an addition of] and/or adding urea, trifluoroethanol or peptides.

16. (Amended) [Process for] A method according to claim 27 further comprising  
increasing the specific physiological activities of molecules by [their incorporation as]  
incorporating a radical R into a compound of the general formula (I) [as defined in any one of  
claims 1 to 9].

Claim 17 has been canceled.

18. (Amended) [Preparation according to claim 17 against] A method of treating  
diseases arising from inflammation, viral and bacterial infections, influenza viruses, selectin-  
 mediated inflammatory processes, tumour metastases, or in the neutralisation of antibodies in  
 autoimmune disorders and transplants; said method comprising administering a compound of  
the general formula (I)



wherein

X is an m-valent unit and

B are identical or different and denote K-R,

wherein

K is a bond or is  $A^1-(A^2-A^3)_k$ -sp, wherein

$A^1$  is  $(CH_2)_t Y(CH_2)_u$ , wherein

Y is  $>C=O$ ,  $>NH$ ,  $-O-$ ,  $-S-$  or a bond,

t is an integer from 0 to 6 and

u is an integer from 0 to 6,

$A^2$  is  $-NHCO-$ ,  $-CONH-$ ,  $-OCONH-$  or  $SCONH-$ , or is  $-CO-$ ,

$A^3$  is  $(CH_2)_r$ ,  $O(CH_2)_r$ ,  $NH(CH_2)_r$ ,  $S(CH_2)_r$  or  $-(CHQ)-$ , wherein

r is an integer from 1 to 6 and

Q is a substituted or unsubstituted alkyl or aryl group,

sp is a divalent spacer or a bond, and



In re: Nikolai Vladimirovich Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 24 of 30

k is an integer from 5 to 100, and  
R is hydrogen; a ligand suitable for specific bonding to a receptor;  
a marker molecule; or a catalytically active group; and  
m is at least 2,  
with the proviso that  
 (1) in the compound at least one R is not hydrogen,  
 (2) there are at least two K that are not a bond, and  
 (3) X, B and m are so selected that an intermolecular association of the K in liquid phase  
by the formation of hydrogen bonds is possible, with formation of aggregates that  
present on the surface a plurality of R that are not hydrogen, and  
 (5) the molar mass of the fragment X(K)<sub>m</sub> is less than 20,000; or  
administering into an aggregate of the general formula (II)  

$$\{X(B)_m\}_n \quad (II)$$

wherein

X(B)<sub>m</sub> may be identical or different and denote a compound of the general  
formula (I), and  
n is from 2 to 100,000,  
and wherein X(B)<sub>m</sub> are non-covalently bonded.

Claim 19 has been canceled.

20. (Amended) [Use of a compound as defined in any one of claims 1 to 9 or of an  
 aggregate as defined in any one of claims 10 to 12 in the preparation of functionalised] A  
method according to claim 18 further comprising preparing functionalized molecular  
surfaces.

Claims 21 and 22 have been canceled.

23. (Amended) [Compound] A compound of the general formula (III),

$$X(B)_m \quad (III)$$

In re: Nikolai Vladimirovich Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 25 of 30

wherein

X is an m-valent unit and

B are identical or different and denote K-H,

wherein

K is  $A^1-(A^2-A^3)_k$ -sp, wherein

$A^1$  is  $(CH_2)_tY(CH_2)_u$ , wherein

Y is  $>C=O$ ,  $>NH$ ,  $-O-$ ,  $-S-$  or a bond,

t is an integer from 0 to 6 and

u is an integer from 0 to 6,

$A^2$  is  $-NHCO-$ ,  $-CONH-$ ,  $-OCONH-$  or  $SCONH-$ , or is  $-CO-$ ,

$A^3$  is  $(CH_2)_r$ ,  $O(CH_2)_r$ ,  $NH(CH_2)_r$ ,  $S(CH_2)_r$  or  $-(CHQ)-$ , wherein

r is an integer from 1 to 6 and

Q is a substituted or unsubstituted alkyl or aryl group,

sp is a divalent spacer or a bond, and

k is an integer from 5 to 100, and

m is at least 2,

with the proviso that

- (1) X, B and m are so selected that an intermolecular association of the K in liquid phase is possible, especially under aqueous conditions, by the formation of hydrogen bonds, with formation of aggregates, and
- (2) the molar mass of the fragment  $X(K)_m$  is less than 20,000, especially less than 4000.

Claim 24 has been canceled.

25. (New) A method of preparing a therapeutic drug comprising:  
 preparing a compound of the general formula (III),



wherein

X is an m-valent unit and

In re: Nikolai Vladimirovich Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 26 of 30

B are identical or different and denote K-H,  
 wherein

K is  $A^1-(A^2-A^3)_k\text{-sp}$ , wherein  
 $A^1$  is  $(\text{CH}_2)_t\text{Y}(\text{CH}_2)_u$ , wherein  
 Y is  $>\text{C}=\text{O}$ ,  $>\text{NH}$ ,  $-\text{O}-$ ,  $-\text{S}-$  or a bond,  
 t is an integer from 0 to 6 and  
 u is an integer from 0 to 6,  
 $A^2$  is  $-\text{NHCO}-$ ,  $-\text{CONH}-$ ,  $-\text{OCONH}-$  or  $\text{SCONH}-$ , or is  $-\text{CO}-$ ,  
 $A^3$  is  $(\text{CH}_2)_r$ ,  $\text{O}(\text{CH}_2)_r$ ,  $\text{NH}(\text{CH}_2)_r$ ,  $\text{S}(\text{CH}_2)_r$  or  $-(\text{CHQ})-$ , wherein  
 r is an integer from 1 to 6 and  
 Q is a substituted or unsubstituted alkyl or aryl group,  
 sp is a divalent spacer or a bond, and  
 k is an integer from 5 to 100, and  
 m is at least 2,

with the proviso that

- (1) X, B and m are so selected that an intermolecular association of the K in liquid phase is possible, especially under aqueous conditions, by the formation of hydrogen bonds, with formation of aggregates, and
- (2) the molar mass of the fragment  $\text{X}(\text{K})_m$  is less than 20,000, especially less than 4000.

26. (New) A method of treating diseases arising from inflammation, viral and bacterial infections, influenza viruses, selectin-mediated inflammatory processes, tumour metastases, or in the neutralisation of antibodies in autoimmune disorders and transplants; said method comprising administering a compound of the general formula (III),



wherein

X is an m-valent unit and  
 B are identical or different and denote K-H,  
 wherein

In re: Nikolai Vladimi h Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 27 of 30

K is  $A^1-(A^2-A^3)_k$ -sp, wherein  
 $A^1$  is  $(CH_2)_tY(CH_2)_u$ , wherein  
 Y is  $>C=O$ ,  $>NH$ ,  $-O-$ ,  $-S-$  or a bond,  
 t is an integer from 0 to 6 and  
 u is an integer from 0 to 6,  
 $A^2$  is  $-NHCO-$ ,  $-CONH-$ ,  $-OCONH-$  or  $SCONH-$ , or is  $-CO-$ ,  
 $A^3$  is  $(CH_2)_r$ ,  $O(CH_2)_r$ ,  $NH(CH_2)_r$ ,  $S(CH_2)_r$  or  $-(CHQ)-$ , wherein  
 r is an integer from 1 to 6 and  
 Q is a substituted or unsubstituted alkyl or aryl group,  
 sp is a divalent spacer or a bond, and  
 k is an integer from 5 to 100, and

m is at least 2,

with the proviso that

- (1) X, B and m are so selected that an intermolecular association of the K in liquid phase is possible, especially under aqueous conditions, by the formation of hydrogen bonds, with formation of aggregates, and
- (2) the molar mass of the fragment  $X(K)_m$  is less than 20,000, especially less than 4000.

27. (New) A method of preparing an aggregate comprising:  
 preparing a compound of the general formula (II)



wherein

$X(B)_m$  may be identical or different and denote a compound of the general formula (I),



wherein

X is an m-valent unit and

B are identical or different and denote K-R,

wherein

K is a bond or is  $A^1-(A^2-A^3)_k$ -sp, wherein  
 $A^1$  is  $(CH_2)_tY(CH_2)_u$ , wherein  
 Y is  $>C=O$ ,  $>NH$ ,  $-O-$ ,  $-S-$  or a bond,

In re: Nikolai Vladimirovich Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 28 of 30

t is an integer from 0 to 6 and  
 u is an integer from 0 to 6,  
 $A^2$  is -NHCO-, -CONH-, -OCONH- or SCONH-, or is -CO-,  
 $A^3$  is  $(CH_2)_r$ ,  $O(CH_2)_r$ ,  $NH(CH_2)_r$ ,  $S(CH_2)_r$  or  $-(CHQ)-$ , wherein  
 r is an integer from 1 to 6 and  
 Q is a substituted or unsubstituted alkyl or aryl group,  
 sp is a divalent spacer or a bond, and  
 k is an integer from 5 to 100, and  
 R is hydrogen; a ligand suitable for specific bonding to a receptor;  
 a marker molecule; or a catalytically active group; and  
 m is at least 2,  
 with the proviso that  
 (1) in the compound at least one R is not hydrogen,  
 (2) there are at least two K that are not a bond, and  
 (3) X, B and m are so selected that an intermolecular association of the K in liquid phase  
 by the formation of hydrogen bonds is possible, with formation of aggregates that  
 present on the surface a plurality of R that are not hydrogen, and  
 (4) the molar mass of the fragment  $X(K)_m$  is less than 20,000, and  
 n is from 2 to 100,000,  
 and wherein  $X(B)_m$  are non-covalently bonded.

28. (New) A method of preparing a therapeutic drug comprising:

preparing the compound of the general formula (I)



wherein

X is an m-valent unit and

B are identical or different and denote K-R,

wherein

K is a bond or is  $A^1-(A^2-A^3)_k-sp$ , wherein

$A^1$  is  $(CH_2)_tY(CH_2)_u$ , wherein

Y is  $>C=O$ ,  $>NH$ , -O-, -S- or a bond,

t is an integer from 0 to 6 and

In re: Nikolai Vladimirovich Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 29 of 30

u is an integer from 0 to 6,  
 $A^2$  is -NHCO-, -CONH-, -OCONH- or SCONH-, or is -CO-,  
 $A^3$  is  $(CH_2)_r$ ,  $O(CH_2)_r$ ,  $NH(CH_2)_r$ ,  $S(CH_2)_r$  or  $-(CHQ)-$ , wherein  
 r is an integer from 1 to 6 and  
 Q is a substituted or unsubstituted alkyl or aryl group,  
 sp is a divalent spacer or a bond, and  
 k is an integer from 5 to 100, and

R is hydrogen; a ligand suitable for specific bonding to a receptor;  
 a marker molecule; or a catalytically active group; and

m is at least 2,

with the proviso that

- (1) in the compound at least one R is not hydrogen,
- (2) there are at least two K that are not a bond, and
- (3) X, B and m are so selected that an intermolecular association of the K in liquid phase by the formation of hydrogen bonds is possible, with formation of aggregates that present on the surface a plurality of R that are not hydrogen, and
- (4) the molar mass of the fragment  $X(K)_m$  is less than 20,000; or  
 preparing the compound of the general formula (II):



wherein

$X(B)_m$  may be identical or different and denote a compound of the general formula (I), and

n is from 2 to 100,000,

and wherein  $X(B)_m$  are non-covalently bonded; and

a pharmaceutically acceptable carrier.

29. (New) A method of preparing a diagnostic test comprising:

providing a test reagent

preparing a compound of the general formula (I)



wherein

In re: Nikolai Vladimi h Bovin et al.  
 Serial No.: To Be Assigned  
 Filed: Concurrently Herewith  
 Page 30 of 30

X is an m-valent unit and

B are identical or different and denote K-R,  
 wherein

K is a bond or is  $A^1-(A^2-A^3)_k\text{-sp}$ , wherein

$A^1$  is  $(\text{CH}_2)_t\text{Y}(\text{CH}_2)_u$ , wherein

Y is  $>\text{C}=\text{O}$ ,  $>\text{NH}$ ,  $-\text{O}-$ ,  $-\text{S}-$  or a bond,

t is an integer from 0 to 6 and

u is an integer from 0 to 6,

$A^2$  is  $-\text{NHCO}-$ ,  $-\text{CONH}-$ ,  $-\text{OCONH}-$  or  $\text{SCONH}-$ , or is  $-\text{CO}-$ ,

$A^3$  is  $(\text{CH}_2)_r$ ,  $\text{O}(\text{CH}_2)_r$ ,  $\text{NH}(\text{CH}_2)_r$ ,  $\text{S}(\text{CH}_2)_r$  or  $-(\text{CHQ})-$ , wherein

r is an integer from 1 to 6 and

Q is a substituted or unsubstituted alkyl or aryl group,

sp is a divalent spacer or a bond, and

k is an integer from 5 to 100, and

R is hydrogen; a ligand suitable for specific bonding to a receptor;  
 a marker molecule; or a catalytically active group; and

m is at least 2,

with the proviso that

- (1) in the compound at least one R is not hydrogen,
  - (2) there are at least two K that are not a bond, and
  - (3) X, B and m are so selected that an intermolecular association of the K in liquid phase by the formation of hydrogen bonds is possible, with formation of aggregates that present on the surface a plurality of R that are not hydrogen, and
  - (5) the molar mass of the fragment  $\text{X}(\text{K})_m$  is less than 20,000; or
- preparing an aggregate of the general formula (II)



wherein

$\text{X}(\text{B})_m$  may be identical or different and denote a compound of the general formula (I), and

n is from 2 to 100,000,

and wherein  $\text{X}(\text{B})_m$  are non-covalently bonded;

and comparing the test reagent to the compounds of the general formula (I) or (II).